

Should ADHD Be Treated If Your Teen Is Using Cannabis?

Ryan Sultan, MD

*What
Parents
Need to
Know*



FOR MANY PARENTS, few situations feel more confusing than watching their teenager struggle with focus, motivation, and organization. While inconsistent motivation and temporary disorganization are common in teenagers as executive function develops, as a parent, it can sometimes be incredibly difficult to discern when behaviors are a clinical issue. This dilemma is further exasperated when you know, or suspect, your teen to be using cannabis. It is natural to wonder whether the attention problems are really ADHD, whether cannabis is causing the symptoms, or whether starting ADHD treatment could somehow make things worse.

ADHD is a neurodevelopmental condition common in children and adolescents characterized by persistent patterns of inattention, impulsivity, and/or hyperactivity that interfere with daily functioning across school, work, and relationships. In a nationally representative sample of teens in the United States, about one in ten adolescents struggle with ADHD. ADHD is associated with approximately a twofold increased risk of substance use disorder in adolescence, and cannabis use is more prevalent among adolescents with ADHD (36.3%) compared to those without ADHD (21.2%). ADHD functions as an early risk amplifier for most substance use issues. The risk mechanism is shown in Figure 1.

ADHD youth have roughly double the risk of cannabis use, with many pathways from early ADHD to substance use. Childhood ADHD symptoms might become behavioral dysregulation in adolescence, leading to chronic failure experiences, emotional instability, and diagnosable mental health conditions.

Given this underlying mechanism, it's understandable that parents may worry that medication could increase addiction risk or inadvertently reinforce substance use. However, using substances like cannabis to self-manage symptoms is fundamentally different from receiving carefully prescribed, evidence-based ADHD treatment. When ADHD is left untreated, individuals are more likely to seek relief through substances, *creating even more risk*. In contrast, appropriate medical treatment targets the underlying condition—reducing impulsivity, improving decision-making, and ultimately lowering the risk of ongoing substance use over time.

The confusion: ADHD, cannabis, or both?

ADHD and cannabis use can overlap in ways that look very similar. A teen with untreated ADHD may struggle with task

initiation, underestimate time, live in a cluttered environment, and begin falling behind academically. A teen who is using cannabis may also look forgetful, slow to start, mentally foggy, and disengaged. Anxiety can add another layer, making concentration even harder and creating a cycle of avoidance, panic, and poor follow-through.

It's natural to think the solution is to stop cannabis use first in order to better understand what's driving motivational and organizational issues. However, this approach can sometimes delay needed support, as ADHD and substance use often interact rather than exist in isolation.

A careful evaluation looks for developmental ADHD markers, not just current symptoms. Were there signs of inattention, disorganization, impulsivity, or time

blindness earlier in childhood, before cannabis entered the picture? Has the teen always struggled with routines, school tasks, or emotional regulation? ADHD is a neurodevelopmental condition. It does not suddenly appear because a teenager started using cannabis. (See figure 2, Developmental Impact of ADHD.)

Why cannabis worsens attention—but does not cause lifelong ADHD

There is certainly evidence that cannabis use can make attention and executive functioning worse. Adolescent cannabis use is associated with slower thoughts, greater difficulty concentrating, poorer academic performance, and broader psychosocial problems. There is an evident dose-related pattern in which more

Figure 1





Figure 2

Developmental Impact of ADHD



frequent cannabis use is associated with worse outcomes.

When a teen is already struggling with attention, school, or mental health, cannabis use can amplify the exact cognitive problems that make ADHD hard to manage. At the same time, cannabis usually does not explain a lifelong history of ADHD symptoms—cannabis may worsen attention, but it rarely accounts for true, longstanding ADHD by itself. Anxiety can further magnify executive dysfunction, which is why teens who use cannabis and feel anxious may appear even more scattered or impaired.

For parents, this means the goal is not to choose one explanation and ignore the others. The goal is to untangle them. Your teen may have ADHD and also be using cannabis. They may be using cannabis partly because they are overwhelmed, anxious, or frustrated by untreated ADHD. Or they may be caught in a loop where

cannabis worsens the very symptoms they are trying to escape.

Why delaying treatment can backfire

A common parental instinct is to postpone ADHD treatment until cannabis use is completely resolved. However, in certain high-risk situations, failing to treat ADHD may be the greatest danger. Why might that be?

Untreated ADHD can make daily life feel unmanageable. Teens with ADHD may struggle with school failure, shame, chronic conflict at home, emotional dysregulation, and a sense that they are always behind. (See figure 3, Emotional Dysregulation and ADHD.) Cannabis can become a way to numb frustration, escape pressure, or quiet internal restlessness. If the underlying ADHD remains untreated, the teen may continue reaching for short-term relief even when

everyone agrees cannabis is making things worse.

This does not mean treatment should be automatic or simplistic. It means that postponing all ADHD care until things are “perfect” may leave the teen stuck in the same cycle. In some families, treating ADHD more effectively becomes part of what helps reduce cannabis reliance, not something that must wait until after the cannabis problem is solved.

A step-by-step approach for parents

When ADHD, cannabis, and anxiety overlap, sequencing—not speed—drives safe care. For many families, the most helpful mindset is this: the goal is not to “win” against your teen’s cannabis use. Rather, the goal is to understand what is happening well enough to build an effective long-term plan. Here is a parent-friendly approach (see also figure 4).

STEP 1: Start with curiosity, not accusation.

If your teen is using cannabis, begin with a collaborative stance. Try to understand what role it is playing. Are they using it socially, to sleep, to calm anxiety, to manage boredom, or to cope with school stress? A teenager is much more likely to open up if they feel you are trying to understand the function of the behavior rather than punish it.

Instead of saying, “You need to stop because this is the problem,” try, “Help me understand what cannabis is doing for you right now.” That shift opens the door to more honest conversations and better clinical information.

STEP 2: Clarify the ADHD story.

Ask whether the attention problems existed before cannabis use. Review school history, teacher feedback, parent observations, and early patterns of forgetfulness, disorganization, or impulsivity. A strong developmental history makes ADHD more likely; a

Figure 3

Emotional Dysregulation and ADHD

Individuals with ADHD are more likely to:



Become overly excited



Focus more on negatives than positives



Develop anxiety or depression




Have relationship problems



Be involved in road rage

Figure 4



sudden change after cannabis use points to a more complicated picture.

STEP 3: Evaluate anxiety and other contributors. Anxiety often makes teens look more inattentive than they are. A mind full of worry cannot focus well. Poor sleep, panic symptoms, depression, and academic burnout can all intensify executive dysfunction. Anxiety must be treated as a major part of the diagnostic picture, not just a side issue.

STEP 4: Consider a brief period of reduced or paused cannabis use to clarify the baseline. This is not always easy, and it may not happen immediately, but even a short stretch of reduced use can help families and clinicians see what symptoms remain when cannabis effects are less active.

STEP 5: Sequence treatment thoughtfully. Once the picture is clearer, treatment can be planned in a safer manner. In some cases, that may mean starting with behavioral supports, CBT, sleep stabilization, anxiety treatment, or parent coaching. In other cases, ADHD medication may be appropriate, especially when symptoms are severe and longstanding. Treatment should be structured and monitored rather than avoided out of fear. It is important to choose approaches that minimize misuse risk and use lower-misuse options when

appropriate at the start.

STEP 6: Think long-term, not just crisis-to-crisis. Parents understandably want immediate answers, but ADHD and substance use are both long-game issues. Joyce Epstein's parenting framework (figure 5) fits especially well here: adolescents respond better to connection, collaboration, and steady scaffolding than to authoritarian control. Teens are more likely to engage in treatment when they feel respected, involved, and supported over time.

This means combining whatever is needed—medication when indicated, CBT or other therapy, motivational work around cannabis, skill-building, school supports, and regular follow-up. The goal is not just stopping one behavior. The goal is helping your teen build a more stable life.

YOU MAY BE THINKING, “I really worry with my family’s substance use history that my child will get really used to taking a substance early in life.” It is important to discuss this concern with the medical professional providing your teen’s treatment.

A family history of substance use does not guarantee that your child will have a substance use

problem. There are a wide range of factors that drive substance use outcomes, not just genetic risk factors (examples include peer influence, trauma and childhood influence, and chronic stress).

When the medical treatment professional responds to the parent, it is first necessary to validate the concern. It is a completely valid concern to not want to teach children to rely solely on substances, pharmacologic or not. We next want to assure parents that pharmacological treatment is only one of the treatments. **A**



A board-certified adult and child and adolescent psychiatrist, **Ryan Sultan, MD**, is an assistant professor of clinical psychiatry at Columbia University College of Physicians and Surgeons, where he runs the Sultan Lab for Mental Health Informatics. He is also the medical director of Integrative Psychiatry of Manhattan. His research and clinical work focus on the intersection of ADHD, adolescent substance use, and mental health, including federally funded studies on cannabis use in youth and the use of AI to improve mental health screening. Learn more about his work at ryansultan.com. He treats patients and trains the next generation of psychiatrists at NewYork-Presbyterian/Columbia.

Figure 5

